

2015

The Effect of Behavioral Interventions on Body Weight in Children: A Systematic Review and Meta-Analysis

Kristina Tymes-Wilbekin

Austin Clark

Kathryn A. Kaiser

Follow this and additional works at: <https://digitalcommons.library.uab.edu/inquiro>



Part of the [Higher Education Commons](#)

Recommended Citation

Tymes-Wilbekin, Kristina; Clark, Austin; and Kaiser, Kathryn A. (2015) "The Effect of Behavioral Interventions on Body Weight in Children: A Systematic Review and Meta-Analysis," *Inquiro, the UAB undergraduate science research journal*: Vol. 2015: No. 9, Article 20.

Available at: <https://digitalcommons.library.uab.edu/inquiro/vol2015/iss9/20>

This content has been accepted for inclusion by an authorized administrator of the UAB Digital Commons, and is provided as a free open access item. All inquiries regarding this item or the UAB Digital Commons should be directed to the [UAB Libraries Office of Scholarly Communication](#).

The Effect of Behavioral Interventions on Body Weight in Children: A Systematic Review and Meta-Analysis

Kristina Tymes-Wilbekin^{1,2,3}, Austin Clark^{4,5}, Kathryn A. Kaiser¹

¹Office of Energetics, School of Public Health, University of Alabama at Birmingham, Birmingham, AL, USA

²PARAdiGM Program, University of Alabama at Birmingham, Birmingham, AL, USA

³Spelman College, Atlanta, GA, USA

⁴Summer in Biomedical Sciences (SIBS) Undergraduate Research Program, University of Alabama at Birmingham, Birmingham, AL, USA

⁵University of West Florida, Pensacola, FL, USA

ABSTRACT

Background: The Centers for Disease Control and Prevention has acknowledged the growing epidemic of childhood obesity and, with it, the likelihood of an increasing percentage of obese adults in the coming years. This problem raises alarm because more people will be susceptible to heart disease, type 2 diabetes and other weight-related illnesses. Thus far, effective interventions have proven elusive.

Methods: We conducted a meta-analysis of pediatric obesity behavioral interventions to examine the effect for improving weight/body composition. We reviewed 14 studies (total N = 3,363) that examined the effects of behavioral therapies on weight change (reduced weight gain while growing or reduced body fat percentage) in children. The studies included are randomized control trials (RCTs) from a prior systematic review that documented weight change as a primary or secondary outcome and were published between January 2007 and July 2009. Using standard meta-analytic methods, we quantified effects to determine whether behavioral therapies are effective in improving body weight/body composition in children.

Results: The overall standardized mean difference of -0.23 with a 95% Confidence Interval of -0.38 to -0.07 indicates that behavioral therapies do have a significant effect as an intervention for improving weight outcomes in children. There was significant heterogeneity present ($I^2 = 61\%$, $p = 0.0003$).

Conclusions: Behavioral interventions may be superior to other types of obesity interventions in children. Larger studies are warranted. These results may be utilized to treat patients, create policies and better inform the public. Future studies might increase intervention duration, which might improve maintenance of effects post-intervention.

Keywords: pediatric obesity, behavioral intervention, weight maintenance, meta-analysis

INTRODUCTION

Permanent weight loss is not an easy process. Most ongoing weight loss interventions are temporary and not effective. This is evident based on the growing obese population. The

Centers for Disease Control and Prevention (CDC) states that childhood obesity has more than doubled in children in the last 30 years¹. Childhood obesity increases the likelihood that a person will be obese later in life, thus contributing to growth of the obese population²⁻⁴.

Obesity has become a major public health concern, and new studies are constantly being published about its prevention and treatment. The most common forms of treatment are diet, exercise, prescription drugs, surgery and behavioral therapies². For this study, behavioral therapies are categorized as those that do not involve surgery or medication and utilize any form of a "lifestyle" education program, cognitive behavioral therapy, parent/guardian involvement, or the creation of support systems. In children, behavioral therapies and lifestyle interventions have shown the best results for weight maintenance³.

A number of studies have reported effects of behavioral therapies for weight maintenance in children, but there have been few meta-analyses quantifying the degree of effects compared to other therapies. After categorizing studies as behavioral, we attempted to analyze whether behavioral therapies have a significant effect on pediatric weight gain and to quantify the effects of this form of intervention versus other forms of treatment. All of the studies evaluated are randomized control trials, which can provide the best measurement of effectiveness of an intervention.

METHODS

The initial step was to collect all the literature that reported behavioral therapy interventions for weight gain prevention in children. These were collected from a previous systematic review⁵.

The following criteria were used in the present review: (1) the collected information was from a human study, (2) the study design used was a RCT, (3) sample sizes were at least 30 before randomization, (4) the intervention lasted 8 weeks or longer, (5) the primary or secondary outcome of the study was weight loss or prevention of weight gain, (6) the article could be acquired in English and (7) the study was published between January 1, 2007 and July 1, 2009. Additionally, studies included had to have utilized a behavioral therapy

intervention. Of the 45 pediatric studies, 15 were categorized as behavioral therapy interventions, 3 of which did not contain the necessary data to be included in the meta-analysis. We attempted to contact the authors for the 3 studies; two responses came after the allotted time, and were later added to the analysis. Of the 14⁶⁻¹⁹ studies included, 5 of the studies had 2 intervention arms^{11, 12, 14, 20}.

For all the studies the raw outcome data extracted was in one of three forms:

- Body Mass Index (BMI)
- Body Weight in kilograms
- z-BMI score (age- and sex-adjusted values of BMI)

To analyze the data, the mean weight or BMI change within all arms of each study was required, along with the standard deviation or standard error. The methods used were adopted from the Cochrane Collaboration Handbook¹⁹. The mean difference and standard deviation for each study were either calculated manually or using Review Manager (RevMan) Software, version 5.3, from the Cochrane Collaboration website²⁰. The RevMan software was also then used to create a forest plot with the included study values. After the initial analysis of data, a subgroup analysis was used to further summarize the data based on three age range subgroups: pre-pubescent (≤ 10 years), early puberty (10–12 years), and post-pubertal (≥ 12 years). In addition, a funnel plot was generated to evaluate the presence of selective bias reporting among the studies (a bias towards studies showing a positive effect).

Finally, a Risk of Bias Assessment was tabulated to gauge different areas of potential bias in each study. The Risk of Bias Assessment was conducted by two authors (KTW and AC) for each individual study. The resulting information was discussed (with KAK) to reach a consensus, and the final results were then tabulated.

RESULTS

The initial systematic review yielded 45 pediatric obesity studies; the selection process is shown in Figure 1. Among the included studies overall, there was a statistically significant effect for behavioral interventions on pediatric weight/body composition. The standardized mean difference (SMD) and confidence interval (CI) are highlighted in Figure 2. Three studies, of the 19 comparisons evaluated, showed results that favored the control versus the treatment arm, meaning there was a negative effect of treatment. The heterogeneity of outcomes among the included studies is significant, $p = 0.0003$ and $I^2 = 61\%$. Figure 2 also displays the SMD of each study and its weight on the overall outcome.

As a post hoc data analysis, a subgroup comparison based on three age ranges was conducted: pre-pubescent (< 10 years), early puberty (10–12 years), and post-pubertal (> 12 years). For the pre-pubescent ($n = 291$) group of studies^{8, 10, 12, 15} the SMD was -0.05 , 95% CI = -0.37 to 0.27 ; heterogeneity of 65%,

$p = 0.009$. In the early puberty ($n = 1,793$) group, the SMD was -0.37 , 95% CI = -0.65 to -0.08 ; heterogeneity of 69%, $p = 0.003$. For the post-pubertal ($n = 1,049$) group^{7, 13, 16, 17} SMD was -0.32 , 95% CI of -0.67 to 0.03 ; heterogeneity = 55%, $p = 0.07$. The funnel plot in Figure 3 indicates that for the studies included in this meta-analysis, there was no evidence of reporting bias. The Risk of Bias Assessment pictured in Figure 4 evaluates potential sources of bias in individual studies as well as the potential risks of bias in the collective group of studies. The most common potential source of bias was the Lack of Attention Placebo in most of the included studies.




Figure 1 | PRISMA diagram of literature search and selection process

DISCUSSION

Our results indicate that behavioral therapies generally have significantly favorable effects on body weight in children. In the post hoc subgroup analysis, by age range/pubertal status, the effect of behavioral therapies on body weight was not significant for the pre-pubescent and post-pubertal age groups. For the early puberty age group, the effect of behavioral therapies did have a significant effect. For the three subgroups, the SMD of effects appear largest in the early puberty age group (10–12 years).




Figure 2 | Forest Plot of Behavioral Therapy Effects on Pediatric Obesity




Figure 3 | Funnel Plot Summary of Reporting Bias

This meta-analysis contained 14 recently published studies ranging in intervention duration from three months to two years. All included studies were randomized control trials, which help to minimize confounding effects. However, this meta-analysis was limited by the small number of studies and the high heterogeneity of observed effects. Future studies should include attention to placebo in their study designs to further minimize the potential for confounds. Additionally, given that other evaluations have indicated that lifestyle interventions are the best intervention for long-term weight maintenance in children⁶, a standardized definition of behavioral therapy is desirable because it would allow better classification and comparison of studies.

In light of the continuing increase of pediatric obesity, the results of this study could be used in support of policies geared toward behavioral therapy-based interventions. However, studies have noted the complexity of designing lifestyle change programs for children because their environments are not isolated¹⁹. The best option for implementing such programs is to include parents in creating goals for their children and incorporating professionals from nutrition, health, and psychological backgrounds in the treatment





Figure 4 | Study-level Risk of Bias (top) and Risk of Bias of all included studies (bottom)

team¹⁹. The more educated the population becomes on ways to reduce weight gain or elicit weight loss in children, the more effective interventions can become in the future.

ACKNOWLEDGMENTS

This research was partially supported by the UAB PARAdiGM Program (NIH/NHLBI R25HL120883-KTW), NIH GM K12GM088010 (PLC), and NIDDK R01DK078826 (DBA). Dr. David Allison was the primary investigator for the systematic review from which included papers were identified. We are grateful to Dr. Allison for his mentorship and use of the data.

REFERENCES

- Childhood Obesity Facts. Centers for Disease Control and Prevention. 2014. Available from: <http://www.cdc.gov/healthyyouth/obesity/facts.htm>.
- Daniels, S.R., Arnett, D.K., Eckel, R.H., Gidding, S.S., Hayman, L.L., Kumanyika, S., et al. Overweight in children and adolescents: pathophysiology, consequences, prevention, and treatment. *Circulation*. 111(15), 1999–2012 (2005).
- Ebbeling, C.B., Pawlak, D.B. & Ludwig, D.S. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 360(9331), 473–82 (2002).
- Freedman, D.S., Khan, L.K., Serdula, M.K., Dietz, W.H., Srinivasan, S.R. & Berenson, G.S. The relation of childhood BMI to adult adiposity: the Bogalusa Heart Study. *Pediatrics*. 115(1), 22–7 (2005).
- Affuso, O., Kaiser, K.A., Carson, T.L., Ingram, K.H., Schwierts, M., Robertson, H., et al. Association of run-in periods with weight loss in obesity randomized controlled trials. *Obesity reviews*. 15(1), 68–73 (2014).
- Wilfley, D.E., Kolko, R.P. & Kass, A.E. Cognitive-behavioral therapy for weight management and eating disorders in children and adolescents. *Child and adolescent psychiatric clinics of North America*. 20(2), 271–85 (2011).
- Doyle, A.C., Goldschmidt, A., Huang, C., Winzelberg, A.J., Taylor, C.B. & Wilfley, D.E. Reduction of overweight and eating disorder symptoms via the Internet in adolescents: a randomized controlled trial. *The Journal of adolescent health*. 43(2), 172–9 (2008).
- Epstein, L.H., Roemmich, J.N., Robinson, J.L., Paluch, R.A., Winiewicz, D.D., Fuerch, J.H., et al. A randomized trial of the effects of reducing television viewing and computer use on body mass index in young children. *Archives of pediatrics & adolescent medicine*. 162(3), 239–45 (2008).
- Foster, G.D., Sherman, S., Borradaile, K.E., Grundy, K.M., Vander Veur, S.S., Nachmani, J., et al. A policy-based school intervention to prevent overweight and obesity. *Pediatrics*. 121(4), e794–802 (2008).
- Golley, R.K., Magarey, A.M., Baur, L.A., Steinbeck, K.S. & Daniels, L.A. Twelve-month effectiveness of a parent-led, family-focused weight-management program for prepubertal children: a randomized, controlled trial. *Pediatrics*. 119(3), 517–25 (2007).
- Janicke, D.M., Sallinen, B.J., Perri, M.G., Lutes, L.D., Huerta, M., Silverstein, J.H., et al. Comparison of parent-only vs family-based interventions for overweight children in underserved rural settings: outcomes from project STORY. *Archives of pediatrics & adolescent medicine*. 162(12), 1119–25 (2008).
- Kalavainen, M.P., Korppi, M.O. & Nuutinen, O.M. Clinical efficacy of group-based treatment for childhood obesity compared with routinely given individual counseling. *International journal of obesity*. 31(10), 1500–8 (2007).
- Kitzman-Ulrich, H., Hampson, R., Wilson, D.K., Presnell, K., Brown, A. & O'Boyle, M. An adolescent weight-loss program integrating family variables reduces energy intake. *Journal of the American Dietetic Association*. 109(3), 491–6 (2009).
- McCallum, Z., Wake, M., Gerner, B., Baur, L.A., Gibbons, K., Gold, L., et al. Outcome data from the LEAP (Live, Eat and Play) trial: a randomized controlled trial of a primary care intervention for childhood overweight/mild obesity. *International journal of obesity*. 31(4), 630–6 (2007).
- Shelton, D., Le Gros, K., Norton, L., Stanton-Cook, S., Morgan, J. & Masterman, P. Randomised controlled trial: A parent-based group education programme for overweight children. *Journal of paediatrics and child health*. 43(12), 799–805 (2007).
- Sichieri, R., Paula Trotte, A., de Souza, R.A., Veiga, G.V. School randomised trial on prevention of excessive weight gain by discouraging students from drinking sodas. *Public health nutrition*. 12(2), 197–202 (2009).
- Tsiros, M.D., Sinn, N., Brennan, L., Coates, A.M., Walkley, J.W., Petkov, J., et al. Cognitive behavioral therapy improves diet and body composition in overweight and obese adolescents. *The American journal of clinical nutrition*. 87(5), 1134–40 (2008).
- Walsh, S.M., Palmer, W., Welsh, J.A. & Vos, M.B. Challenges and successes of a multidisciplinary pediatric obesity treatment program. *Nutrition in clinical practice*. 29(8), 780–785 (2014).
- Wilfley, D.E., Stein, R.I., Saelens, B.E., Mockus, D.S., Matt, G.E., Hayden-Wade, H.A., et al. Efficacy of maintenance treatment approaches for childhood overweight: a randomized controlled trial. *JAMA*. 298(14), 1661–73 (2007).
- Cochrane Handbook for Systematic Reviews of Interventions: The Cochrane Collaboration. 2009. Available from: www.cochrane-handbook.org.
- Collaboration. The Cochrane Collaboration. Review Manager Version 5.3. Copenhagen: The Nordic Cochrane Centre. 2012.